Project:	**Annual ryegrass control in wheat**	
Trial:	**AM1318 - Disc planted**	
District:	**Mullaley**	
Variety:	**Crusader**	
Planting date:	**20/06/2013**	
Application dates:	**20/06/2013**	**IBS**
	20/06/2013	**PSPE**
Soil type:	**Black vertosol**	
Planting Depth:	**3.5-4.5cm**	

Disclaimer:

This document is based on the results from an individual trial and may contain experimental use patterns that are currently off-label. **This document does not provide any interpretation and should not be taken as an endorsement of any unregistered use pattern.**

Professional advice should be sought for specific recommendations to ensure access to the most up to date information and knowledge.

Any product referred to in this document must be used strictly as directed, and in accordance with all label or permit instructions. Always consult the label prior to use.

Disc Planted (Black vertosol)				11/7/2013 (21 DAP)	27/8/2013 (68DAP)	22/9/2013 (94DAP)
Trt No.	**Product**	**Rate ml or g/ha**	**Timing**	**Wheat Establishment Plants/m^2**	**ARG Plants/m^2**	**ARG Plants/m^2**
1	Untreated	-	-	108 a	5.8 a	6.7 a
2	Boxer Gold	2500	IBS	30 c	0.1 c	0.0 b
3	Boxer Gold	2500	PSPE	64 b	0.0 c	0.0 b
5	TriflurX	1500	IBS	20 c	0.9 bc	0.9 b
6	Sakura	118	IBS	67 b	0.1 c	0.2 b
7	Stomp Xtra	1400	IBS	63 b	0.8 bc	0.4 b
8	Boxer Gold + TriflurX	2500 + 800	IBS	24 c	0.1 c	0.2 b
9	Boxer Gold + TriflurX	2500 + 1500	IBS	16 c	0.2 c	0.2 b
10	Boxer Gold + Stomp Xtra	2500 + 1400	IBS	25 c	0.1 c	0.1 b
			P =	<0.01	<0.01	<0.01
			LSD =	14.6	Arcsin(sqrt(x/100)) transformation	Arcsin(sqrt(x/100)) transformation