

Project: Windmill grass control in chickpeas
Trial: AM1311
District: Baradine
Variety: PBA HatTrick
Planting date: 21/05/2013
Planter type: Disc planter

Disclaimer:

This document is based on the results from an individual trial and may contain experimental use patterns that are currently off-label. **This document does not provide any interpretation and should not be taken as an endorsement of any unregistered use pattern.**

Professional advice should be sought for specific recommendations to ensure access to the most up to date information and knowledge.

Any product referred to in this document must be used strictly as directed, and in accordance with all label or permit instructions. Always consult the label prior to use.

Application dates:

20/05/2013	IBS
21/05/2013	PSPE

Trt No.	Product	Rate ml or g/ha	Timing	16/11/2013 179 DAP		7/04/2014 322 DAP	
				% of Plot covered by weed		Weeds/m ²	
				Wireweed*	Medicago spp	Grasses	
1	Untreated	-	-	10	43 b	169 a	
2	Sakura	118	IBS	25	48 ab	88 cd	
3	Outlook	1000	IBS	40	48 ab	143 ab	
4	Avadex Xtra	1600	IBS	23	65 ab	104 bc	
5	Triflur Xcel	1700	IBS	0	55 ab	133 ab	
6	Stomp Xtra	2250	IBS	0	55 ab	48 d	
7	Balance	100	PSPE	18	0 c	135 ab	
8	Terbyne	1400	PSPE	0	16 c	144 ab	
9	Balance + Simazine	100 + 1500	PSPE	3	0 c	113 bc	
10	Gp B S	25	PSPE	0	46 b	84 cd	
11	Terbyne + Gp B S	1000 + 20	PSPE	0	73 a	114 bc	
12	Terbyne + Gp B S + Balance	1000 + 20 + 80	PSPE	0	0 c	120 bc	
				P =	0.16	0.00	<0.01
				LSD =	nsd	25.8	44.7

*Wireweed unevenly distributed across the site

Field comments:

26/09/2013 Trial sprayed out with 1.8L/ha Roundup Attack + 1.8L/ha Amicide 625 + wetter

16/11/2013 Drought conditions. Windmill grass either dead or dormant.

Difficult to assess due to the dried state of weed species

7/04/2014 Large germination of grass weeds. Estimate 80% ABYG and 20% WMG