

Project: Residual Herbicides for Control of Chloris spp.
Trial: BB1204
Location: Nandi-St Ruth
Soil Type: Sandy Loam
Application Timing: Pre-emergent
Application Date: 6/12/2012
Equipment: 4m Quad-bike mounted boom
Nozzles: AIXR110015
Nozzle Pressure (kPa): 300
Speed (km/hr): 10.3
Volume (L/ha): 70
Target: Feathertop Rhodes grass (*Chloris virgata*)

Disclaimer:

This document is based on the results from an individual trial and may contain experimental use patterns that are currently off-label. **This document does not provide any interpretation and should not be taken as an endorsement of any unregistered use pattern.**

Professional advice should be sought for specific recommendations to ensure access to the most up to date information and knowledge.

Any product referred to in this document must be used strictly as directed, and in accordance with all label or permit instructions. Always consult the label prior to use.

Trt No.	Treatment	Rate ml or g/ha	Feathertop Rhodes Grass (<i>Chloris virgata</i>)
			6/02/2013 62 DAA FTR /m ²
1	Untreated	-	72.3 a
2	Flame	150	33.7 abcd
3	Flame	200	15.3 def
4	Flame + Gp C D	150 + 1000	11.6 defg
5	Flame + Balance	150 + 100	2.6 fg
6	Flame + Terbyne	150 + 1000	21.7 bcdef
7	Flame + Nutrazine	150 + 1000	11.9 defg
8	Stomp Xtra	4500	0.1 g
9	Gp K S	118	5.3 efg
10	Glean + Balance	20 + 100	18.9 cdef
11	Primextra Gold	3200	27.6 bcde
12	Boxer Gold	2500	57.8 ab
13	Dual Gold	2000	9.1 defg
14	Balance	100	14.0 def
15	Terbyne	1000	55.4 abc
16	Gp G V	180	3.9 fg
		P =	<0.01
		LSD =	Arcsin Transformation

Treatment means followed by the same letter are not significantly different at P = 0.05

DAA= Days after application