

Project: Fungicides on Sorghum
Trial: BB1301
District: Millmerran
Paddock History: Sorghum
Crop: Sorghum
Rowing Spacing (cm): 100cm
Sowing Depth (mm): 40mm
Soil Type: Loam
Application Timing: Mid flowering
Application Date: 13/03/2013
Equipment: 4m Quad-bike mounted boom
Nozzles: AIXR110015
Nozzle Pressure (kPa): 300
Speed (km/hr): 10.3
Volume (L/ha): 70
Target: Leaf Blight like symptoms (not positively identified)

Disclaimer:

This document is based on the results from an individual trial and may contain experimental use patterns that are currently off-label. **This document does not provide any interpretation and should not be taken as an endorsement of any unregistered use pattern.**

Professional advice should be sought for specific recommendations to ensure access to the most up to date information and knowledge.

Any product referred to in this document must be used strictly as directed, and in accordance with all label or permit instructions. Always consult the label prior to use.

Trt No.	Treatment	Rate ml or g/ha	Adjuvant	Timing	Sorghum
					24/04/2013 42 DAA Incidence of 'Leaf Blight' % of top 5 leaves infected
1	Untreated	-	-	-	26
2	Gp 3/11 AX	200	2% Adigor	50% flowering	22
3	Gp 3/11 AX	400	2% Adigor	50% flowering	25
4	Gp 3/11 AX	800	2% Adigor	50% flowering	21
5	Gp 3 Pr	300	-	50% flowering	25
6	Gp 3 F	290	-	50% flowering	20
7	Gp 3 TX	500	-	50% flowering	26
8	Gp 3/11 Op	1000	-	50% flowering	21
9	Gp 3 Op	500	-	50% flowering	23
				P =	0.49
				LSD =	nsd

Treatment means followed by the same letter are not significantly different at P = 0.05

nsd= no significant difference

DAA= days after application

No indication of any fungicide activity on 'Leaf Blight' in this trial