

Project: Impact of Sorghum Hybrids on *Pratylenchus thornei*
Trials: BB1210-1217, BB1317-1318
Districts: Darling Downs to south of Moree

Disclaimer:

This document is based on the results from an individual trial and may contain experimental use patterns that are currently off-label. **This document does not provide any interpretation and should not be taken as an endorsement of any unregistered use pattern.**

Professional advice should be sought for specific recommendations to ensure access to the most up to date information and knowledge.


Any product referred to in this document must be used strictly as directed, and in accordance with all label or permit instructions. Always consult the label prior to use.

1. Comparison of Sorghum hybrids evaluated in 9 common trials

Hybrid	<i>Pratylenchus thornei</i> Near' Harvest <i>Pt</i> /g soil 8 cores/plot @ 0-30cm
MR-Bazley	6.2
MR-Buster	5.5
MR-Eclipse	6.3
Pacific MR 43	6.3
Pioneer G22	5.1
Pioneer G33	6.3
P=	0.77
LSD=	nsd

No indication of any difference in resistance level to *Pt* between Sorghum hybrids

2. Comparison of all evaluated Sorghum hybrids relative to MR-Buster



10 trials over 2 seasons

No significant difference between any hybrid, in any trial

No indication of any difference in resistance levels to *Pt* between Sorghum hybrids